


**Precalculus**  
**Lesson 9.4: Vectors**  
**Mrs. Snow, Instructor**

Many concepts in science involve applications of mathematics that measure certain quantities by their magnitude like length, mass, area, temperature, or energy. Only one number is needed to describe a length of 7 inches or 5°C for example. This single quantity is called **scalar**.

There are, however, many applications that involve not only the *magnitude* of an object but also, the *direction* of the displacement.

**vector**: a quantity that has both magnitude and direction. For example, the flight pattern of a plane, has both *speed (magnitude)* and *direction* of travel. Velocity, acceleration, and force are described by both magnitude and direction and are known as vectors.



P is the initial point  
 Q is the terminal point

All vectors have two things:

**Direction** – follow the arrow.


**Magnitude** – the length of the vector.

**Graphing Vectors**


Use the vector to graph each of the following vectors:



$v - w$



$2v + 3w$



## Find a Position Vector

If we locate a vector in a coordinate plane we can describe it analytically by writing it in components.


Vector  $\mathbf{v}$ , may be described with initial point  $P_1(x_1, y_1)$  and terminal point  $P_2(x_2, y_2)$ , therefore:

$$\mathbf{v} = \langle x_2 - x_1, \quad y_2 - y_1 \rangle$$

$$\mathbf{v} = \langle \mathbf{a}, \mathbf{b} \rangle$$

This vector may be called the **position vector or component form**

$$\mathbf{v} = \langle a, b \rangle = \langle x_2 - x_1, y_2 - y_1 \rangle$$



Find the position vector  $\mathbf{v}$  with initial point  $(-1, 2)$  and terminal point  $(4, 6)$ .

## Vectors in terms of $\mathbf{i}$ and $\mathbf{j}$

A vector of length  $1$  is called a **unit vector**. The vector  $\mathbf{w} \left\langle \frac{3}{5}, \frac{4}{5} \right\rangle$  is an example of a **unit vector**. We have two special unit vectors  $\mathbf{i}$  and  $\mathbf{j}$ .

“ $\mathbf{i}$ ” is a unit vector in the x-direction and “ $\mathbf{j}$ ” is a unit vector in the y-direction. Any vector in the x-direction can be written as a scalar multiple of  $\mathbf{i}$  and any vector in the y-direction can be written as a scalar multiple of  $\mathbf{j}$ . They are defined as:

$$\mathbf{i} = \langle 1, 0 \rangle \text{ and } \mathbf{j} = \langle 0, 1 \rangle, \text{ where } \|\mathbf{i}\| = \sqrt{1^2 + 0^2} \text{ and } \|\mathbf{j}\| = \sqrt{0^2 + 1^2}.$$

Any vector may be express in terms of **i** and **j**.

### Algebraic Operations

Vectors may be added, subtracted, or have scalar multiplication. Pretty straight forward:

Let  $\mathbf{v} = a_1\mathbf{i} + b_1\mathbf{j} = \langle a_1, b_1 \rangle$  and  $\mathbf{w} = a_2\mathbf{i} + b_2\mathbf{j} = \langle a_2, b_2 \rangle$  be two vectors, and let  $\alpha$  be a scalar. Then

$$\mathbf{v} + \mathbf{w} = (a_1 + a_2)\mathbf{i} + (b_1 + b_2)\mathbf{j} = \langle a_1 + a_2, b_1 + b_2 \rangle \quad (2)$$

$$\mathbf{v} - \mathbf{w} = (a_1 - a_2)\mathbf{i} + (b_1 - b_2)\mathbf{j} = \langle a_1 - a_2, b_1 - b_2 \rangle \quad (3)$$

$$\alpha\mathbf{v} = (\alpha a_1)\mathbf{i} + (\alpha b_1)\mathbf{j} = \langle \alpha a_1, \alpha b_1 \rangle \quad (4)$$

$$\|\mathbf{v}\| = \sqrt{a_1^2 + b_1^2} \quad (5)$$

If  $v = 2i + 3j = \langle 2, 3 \rangle$  and  $w = 3i - 4j = \langle 3, -4 \rangle$ ,

find: a)  $v + w$ ,    b)  $v - w$ ,    c)  $3v$ ,    d)  $2v - 3w$ ,    and  $\|v\|$

A vector that represents speed and velocity of an object is called a **velocity vector**. A vector describing a force represents the direction and amount of force acting upon an object and is called a **force vector**.

### Find a Vector from its Direction and Magnitude

Given the magnitude  $\|v\|$  of a nonzero vector  $\mathbf{v}$  and the **direction angle**  $\alpha$ ,  $0^\circ < \alpha < 360^\circ$ , between  $\mathbf{v}$  and  $\mathbf{i}$ , then:

$$v = \|v\|(\cos \alpha \mathbf{i} + \sin \alpha \mathbf{j})$$

### Writing a Vector When Its Magnitude and Direction Are Given

A ball is thrown with an initial speed of 25 mph in a direction that makes an angle of  $30^\circ$  with the positive x-axis. Express the velocity vector  $\mathbf{v}$  in terms of  $\mathbf{i}$  and  $\mathbf{j}$ . What is the initial speed in the horizontal direction? What is the initial speed in the vertical direction?

### Finding the Direction Angle of a Vector

Find the direction angle  $\alpha$  for  $v = 4\mathbf{i} - 4\mathbf{j}$

### Finding the Actual Speed and Direction of an Aircraft

A Boeing 737 aircraft maintains a constant airspeed of 500 mph headed due south. The jet stream is 80 mph in the northeasterly direction.

- a) Express the velocity  $v_a$  of the 737 relative to the air and velocity  $v_w$  of the jet stream in terms of  $i$  and  $j$ .
- b) Find the velocity of the 737 relative to the ground.
- c) Find the actual speed and direction of the 737 relative to the ground.

### Finding the Weight of a Piano

Two movers require a magnitude of force of 300 pounds to push a piano up a ramp inclined at an angle  $20^\circ$  from the horizontal. How much does the piano weigh?

An Object in Static Equilibrium: the object is at rest and the sum of all forces acting on the object is zero, a.k.a. the resultant force is zero.

A box of supplies that weighs 1200 pounds is suspended by two cables attached to the ceiling. What are the tensions in the two cables?

