Algebra II

Lesson 8-3: Logarithmic Functions as Inverses Mrs. Snow. Instructor

Logarithms are used to express very large numbers or very small numbers in a more manageable way. The common logarithm, base 10, is used in science and engineering applications. Base 2 is used in computer science applications and base e, the natural logarithm, is used in the natural sciences and in pure mathematics such as calculus. Remember our problems where we had to guess and check to figure out the numbers of years it would take to save a certain amount of money? Well, a logarithm will undo that exponent and allow us to easily and quickly solve for the exponent.

Recall that each algebraic operation has an opposite or an inverse:

Operation	Inverse	
multiplication	division	
addition	subtraction	
square: x^2	square root: √	
cube: x^3	cube root: ³ √	

Logarithms are the "opposite" (inverse) of exponents, just as subtraction is the opposite of addition. Logs "undo" exponentials. Technically speaking, logs are the inverses of exponentials. More practically speaking, you can think of logs in terms of the relationship:

To change forms always start with the base. The base is the base regardless of it being a log or an exponential.

Since exponential and logs are inverses, when we have equation in one form, we can write it in the other form. The **common logarithm** is base 10 and unless otherwise written, log is understood to be base 10. So: $log_{10}y = log y$.

Write $10^2 = 100$ in logarithmic form $10^2 = 100 \text{ and } \log_{10} 100 = 2$	$49 = 7^{2}$ $105 - 49 = 2$	base = 7
$10^{-1} = \frac{1}{10}$ Wase 10	729 = 36 $6 = 729$	base = 3

If $2^x = 2^4$. what does x equal??????

We can also go from log to exponent thus solving for x:

Evaluate

$$log_8 16 = x$$

$$8 = 16$$

$$2 = 24$$

$$2 = 24$$

$$2 = 4$$

$$3 = 4$$

- 1. convert to exponential form
- 2. write each side as 2 to a power
- 3. power property of exponents
- 4. set the exponents equal to each other
- 5. solve for x

$$\log_4 2 = x$$

$$4^{\gamma} = 2 \qquad 2 \times 1$$

$$2^{20} = 2^{0} \quad \chi = \frac{1}{2}$$

$$\chi = \frac{1}{2}$$

$$log_4 4 = x$$

$$log_6 1 = x$$

$$log_{10}0 = x$$

no solution

The last examples show us that:

$$log_b b = 1$$


$$log_h 1 = 0$$

 $log_b b = 1$ $log_b 1 = 0$ $log_b 0$ is undefined for all base

Natural Logarithm: The inverse of the function $y = e^x$ is $log_e y = x$ or ln y = x. remember: $log_e = ln$

When a relationships in the real world needs to be described with a logarithmic function, they are best described using the natural logarithms.

Evaluate: $ln e^6 = (o$



Write in exponential form:

Write in exponential form:

$$7.389 = e^x$$

Write in exponential form:

$$ln 5 = 1.609$$

Chemistry While in this section, we will cover pH at the end of this chapter pH of a substance is the measure of acidity of a liquid which is related to the hydrogen ion concentration ($[h^+]$) of the liquid. A liquid becomes more acidic as the number of hydrogen ions increases. A logarithmic equation relates pH to concentration hydrogen ions:

$$pH = -log[h^+]$$

Optional:

Find the concentration of hydrogen ions, [H⁺] lime juice pH = 2.2 2.2 = $\bigcirc log_{10}H^+ \longrightarrow -2.2 = los_{10}H^+$ 10 - 2.2 = $H^+ = .00 (63)$

- log is understood to be base 10
- rewrite with base 10 written and move negative to other side.
 convert into exponential form solve


cider vinegar pH=3.1 egg white pH=8.0
$$8.0 = -l\omega_3$$
, H^+
 $3.1 = -l\omega_3$, H^+ $10^{-3.1} = H^+$ $-8.0 = l\omega_3$, H^+
 $-3.1 = l\omega_3$, O^{-4} $= 1.943 \times 10^{-4}$ $= 1 \times 10^{-8}$

Find the inverse of each logarithmic function:

1. Write in exponential form,

2. Switch: x is y and y is x.

move



Since we really don't have the capabilities to calculate y values for a log base b, we need to look at the inverse of a log function. The inverse of a log is an exponential function.

Graph: $log_5 x = y$

1. rewrite in exponential form:

$$5^9 = x$$


2. By the definition of logarithm, the log is the inverse (x is y and y is x).

3. graph the exponential function

	nential:	$log_5 x = log: Su$	114581
X	У	学二X	6 = Y
-2	25	1 25	-2
-1	15	15	-1
0	1	1	0
1	5	5	1
2	25	25	2

(reflection across line of y = x)

4. reverse the coordinates for the x and y values and plot $y = log_5 x$



When the log function gets more complicated, you will need to use transformations

Parent function	$y = log_b x$	
stretch a>1	$y = alog_b x$	
shrink 0 <a<1< td=""><td>big a the curve stretches or gets larger, vs. fraction a curve gets very thin</td></a<1<>	big a the curve stretches or gets larger, vs. fraction a curve gets very thin	
shift up/down	$y = log_b(x - h) + k$	
shift left/right	where the h is a horizontal shift and the k is our vertical shift	

